

Tubular Field Theory

Izawa K.-I.

*Department of Physics and RESCEU, University of Tokyo,
Tokyo 113-0033, Japan*

Abstract

We propose tubular field theory, which is a continuum analogue of lattice field theory. One-dimensional links (and zero-dimensional sites) in lattice field theory are replaced by two-dimensional tubes to result in two-dimensional spacetime microscopically. As an example, scalar and gauge fields are considered in ‘three-dimensional’ tubular spacetime.

It is conceivable that quantum gravity effects yield microscopic fluctuations of spacetime such as spacetime foam at small scales. The observed four-dimensional spacetime might be macroscopic coarse-graining of such microscopic structures.

In this paper, we consider two-dimensional surface as a simplest tractable model of the microscopic structure and macroscopic spacetime manifold to be large-scale approximation thereof. Although the motivation is closely related to gravity, we restrict ourselves to non-gravitational (and ‘three-dimensional’) setup as a first step.

For concreteness, let us imagine the three-dimensional cubic lattice with its one-dimensional links replaced by two-dimensional cubic tubes of the size l (the area of the cubic tube T with topology $S^1 \times [0, l]$ amounts to $4l^2$ since the circumference of the square S^1 is $4l$). At the ‘site’ $(i, j, k) \in \mathbf{Z}^3$, six cubic tubes are joined with two tubes extended in the two opposite directions along each axis x, y , or z . These tubes constitute a two-dimensional surface Σ as a whole and microscopically the spacetime is two-dimensional.

As a first example, we consider a scalar field on the surface Σ with the action

$$S = \int_{\Sigma} d^2\xi \left(\frac{1}{2} \partial_a \phi \partial^a \phi - V(\phi) \right), \quad (1)$$

where $a = 1, 2$ denotes a two-dimensional index.

At large scales, dropping the Kaluza-Klein modes of ϕ in the circular direction S^1 of the tube $T \sim S^1 \times [0, l]$, we obtain

$$S \simeq \sum_{(i,j,k)} \sum_{\mu} \int_{T_{\mu}} d^2\xi \left(\frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - V(\phi) \right), \quad (2)$$

where $\mu = x, y, z$ represents the positive direction along each axis perpendicular to the circular direction and T_{μ} denotes the corresponding tube at the ‘site’ (i, j, k) . Further approximation goes as follows à la lattice field theory:

$$S \simeq 4l^2 \sum_{(i,j,k)} \left(\frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi - 3V(\phi) \right) \simeq \int dx dy dz \left(\frac{1}{2} \partial_{\mu} \tilde{\phi} \partial^{\mu} \tilde{\phi} - \tilde{V}(\tilde{\phi}) \right), \quad (3)$$

where we have defined

$$\tilde{\phi} = \frac{1}{\sqrt{2}l} \phi, \quad \tilde{V}(\tilde{\phi}) = \frac{3}{2l} V(\phi). \quad (4)$$

In fact, coarse-graining through path integration affects the form of $\tilde{V}(\tilde{\phi})$ and relevant terms in it should become dominant.

As a next example, we consider (nonabelian) gauge theory on Σ with gauge field A_a and charged scalar ϕ . The gauge-invariant action is given by

$$S = \int_{\Sigma} d^2\xi \left(\frac{1}{4} F_{ab} F^{ab} + D_a \phi (D^a \phi)^* - V(\phi) \right). \quad (5)$$

Let us adopt an axial gauge where the gauge field component along the circular direction is vanishing $A_{S^1} = 0$. Dropping the Kaluza-Klein modes, as is the case for the scalar field above, we obtain

$$S \simeq \sum_{(i,j,k)} \sum_{\mu} \int_{T_{\mu}} d^2\xi (D_{\mu} \phi (D^{\mu} \phi)^* - V(\phi)). \quad (6)$$

Further approximation goes as follows:

$$S \simeq 4l^2 \sum_{(i,j,k)} (D_{\mu} \phi (D^{\mu} \phi)^* - 3V(\phi)) \simeq \int dx dy dz (D_{\mu} \tilde{\phi} (D^{\mu} \tilde{\phi})^* - \tilde{V}(\tilde{\phi})), \quad (7)$$

where we have defined

$$\tilde{\phi} = \frac{1}{\sqrt{2}l} \phi, \quad \tilde{V}(\tilde{\phi}) = \frac{3}{2l} V(\phi). \quad (8)$$

In this case, coarse-graining through path integration should generate the kinetic term for the gauge field A_{μ} with gauge invariance, which remains after ‘higher-dimensional’ gauge fixing $A_{S^1} = 0$.

The above two examples imply that three-dimensional (and more generally, higher-dimensional) field theories may arise as large-scale approximations even though microscopically the underlying spacetime is two-dimensional (or more generally, lower-dimensional). To make this approach more realistic, it seems necessary to include (chiral) fermions and gravity in the present framework. In particular, introduction of gravity may naturally make effective dimensions of the macroscopic spacetime to be variable, which should be determined dynamically. Since the underlying spacetime is two-dimensional rather than one-dimensional, such higher-dimensional geometry may be accommodated intrinsically.

Acknowledgments

We would like to thank T. Watari and T. Yanagida for valuable discussions.